

ARIC Manuscript Proposal # 1295

PC Reviewed: 10/09/07
SC Reviewed: _____

Status: A
Status: _____

Priority: 2
Priority: _____

1.a. Full Title: Association of Chronic Obstructive Pulmonary Disease with Venous Thromboembolism in the Atherosclerosis Risk in Communities (ARIC) Study

b. Abbreviated Title (Length 26 characters): COPD and VTE in ARIC

2. Writing Group:

Writing group members: Alanna Chamberlain, MPH
Aaron Folsom, MD
Mary Cushman, MD
Wayne Rosamond, PhD
Susan Heckbert, MD, PhD
Neil Zakai, MD

I, the first author, confirm that all the coauthors have given their approval for this manuscript proposal. AMC **[please confirm with your initials electronically or in writing]**

First author: Alanna Chamberlain
Address: 1300 S. 2nd Street, Suite 300
Minneapolis, MN 55454

Phone: 612-625-5352 Fax: 612-624-0315
E-mail: gram0119@umn.edu

Corresponding/senior author (if different from first author correspondence will be sent to both the first author & the corresponding author):

Address: Aaron Folsom
1300 S. 2nd Street, Suite 300
Minneapolis, MN 55454

Phone: 612-626-8862 Fax: 612-624-0315
E-mail: folsom@epi.umn.edu

3. Timeline:

Statistical Analysis:	September 2007 – October 2007
Manuscript Preparation:	October 2007 – November 2007
Manuscript Revision:	November 2007
Manuscript Submission:	December 2007

4. Rationale:

Chronic obstructive pulmonary disease (COPD) encompasses chronic bronchitis and emphysema, and is the fifth leading cause of death worldwide.¹ Symptoms of COPD include wheezing, dyspnea, sputum production, airflow obstruction, decreased expiratory flow, loss of lung elasticity, hyperinflation, and inflammatory narrowing of airways due to infiltration by neutrophils, macrophages, and CD8-positive T cells.^{1,2,3}

Patients hospitalized for acute exacerbations of COPD may be at risk for developing venous thromboembolism (VTE) because of the presence of bronchial infection and decreased mobility.⁴ VTE comprises deep vein thrombosis (DVT) and pulmonary embolism (PE). Risk factors for VTE include prolonged immobility, trauma, surgery, cancer, previous VTE, increasing age, estrogen use, and congenital and acquired thrombophilic disorders.^{4,5}

Few studies have shown an association between COPD and incidence of VTE. A review article reported that approximately 10% of patients with acute exacerbations of COPD suffer from DVT. The prevalence of PE has been mostly reported in autopsy series where up to 30% of cases who died from acute exacerbations of COPD had PE.⁴ A study of 56 patients, 46 men and 10 women, found DVT and/or PE in 9 (~16%) cases. The investigators of this study did not find a relationship between DVT and/or PE with pulmonary function, arterial blood gas, height, weight, or disease duration,⁶ which suggests decreased mobility from COPD may have contributed to DVT and PE in these patients.

The limited number of studies that have investigated the association between COPD and VTE, as well as a lack of information from large prospective studies, warrants further examination of the association between COPD and incident VTE. Therefore, we propose to examine the relationship between COPD and incident VTE in the ARIC cohort.

References:

1. Pauwels PRA, Rabe KF. Burden and clinical features of chronic obstructive pulmonary disease (COPD). *The Lancet*. 2004/8/14;364:613-620.
2. De Palo VA. Pulmonary disease: Pneumonia, chronic obstructive pulmonary disease, asthma, and thromboembolic disease. *J Am Podiatr Med Assoc*. 2004;94:157-167.
3. Snoeck-Stroband JB, Postma DS, Lapperre TS, et al. Airway inflammation contributes to health status in COPD: A cross-sectional study. *Respir Res*. 2006;7:140.
4. Ambrosetti M, Ageno W, Spanevello A, Salerno M, Pedretti RFE. Prevalence and prevention of venous thromboembolism in patients with acute exacerbations of COPD. *Thrombosis Research*. 2003;112:203-207.
5. Tapson VF. The role of smoking in coagulation and thromboembolism in chronic obstructive pulmonary disease. *Proc Am Thorac Soc*. 2005;2:71-77.
6. ERELEL M, ÇUHADARO G Ç, ECE T, ARSEVEN O. The frequency of deep venous thrombosis and pulmonary embolus in acute exacerbation of chronic obstructive pulmonary disease. *Respiratory Medicine*. 2002/7;96:515-518.

5. Main Hypothesis/Study Questions:

We hypothesize that the incidence of VTE among participants in the ARIC cohort will be significantly higher among those with prevalent COPD compared to those without COPD.

6. Design and analysis (study design, inclusion/exclusion, outcome and other variables of interest with specific reference to the time of their collection, summary of data analysis, and any anticipated methodologic limitations or challenges if present).

Participants from ARIC who have no history of VTE or warfarin use at baseline will be included in this study. The independent variable will be prevalence of COPD as determined by chronic symptoms (cough and sputum production) and lung function tests (FEV₁ as a percentage of predicted value, FVC as a percentage of predicted value, and FEV₁/FVC ratio). The Global Initiative for Chronic Obstructive Lung Disease (GOLD) criteria⁷ will be used to define prevalent COPD in five categories: none, stage 0, mild, moderate, and severe as shown below.

No COPD	FEV ₁ /FVC > 70% FEV ₁ > 80% No chronic symptoms
Stage 0 COPD	FEV ₁ /FVC > 70% FEV ₁ > 80% Chronic symptoms (cough and/or sputum production)
Mild COPD	FEV ₁ /FVC < 70% FEV ₁ ≥ 80% With or without chronic symptoms
Moderate COPD	FEV ₁ /FVC < 70% FEV ₁ < 80% With or without chronic symptoms
Severe COPD	FEV ₁ /FVC < 70% FEV ₁ < 30%

Baseline data for chronic symptoms and lung function will be used for participants who have a VTE event between baseline and visit 2. For the remaining participants, lung function will be averaged over baseline and visit 2, and cough and/or sputum production at either one or both visits will be used to define presence of chronic symptoms. Also, we may combine moderate and severe COPD into a single category depending on the number of participants with moderate and severe COPD. Additionally, if there are not adequate numbers in the COPD categories, we may use quartiles of FEV₁ and FVC as the

independent variables. The dependent variable in this study will be incidence of VTE; however, those who have a cancer-related VTE event will be excluded from the analysis. Cox proportional hazards regression will be used to determine the hazard ratios of VTE by prevalence and severity of COPD, as well as quartiles of FEV₁ and FVC if necessary. If confounding is observed, the associations will be adjusted for the following covariates at baseline: age, sex, race, body mass index, diabetes, factor VIII and smoking status and amount. An interaction test by sex will be conducted, and analyses will be reported separately if evidence of heterogeneity by sex is present. We will also describe the percentage of VTE hospitalizations among the COPD exposure categories (COPD and no COPD at baseline/visit 2) who also had a COPD discharge diagnosis during their VTE hospitalization. Finally, we will examine the main effect of smoking on VTE and test an interaction between smoking and COPD on incidence of VTE. If the smoking-COPD interaction is significant, we will report analyses stratified by smoking status or categories of pack-years of smoking.

7. Pauwels RA, Buist AS, Calverley PMA, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO global initiative for chronic obstructive lung disease (GOLD) workshop summary. *Am J Respir Crit Care Med.* 2001;163:1256-1276.

7.a. Will the data be used for non-CVD analysis in this manuscript? Yes
 No

b. If Yes, is the author aware that the file ICTDER02 must be used to exclude persons with a value RES_OTH = "CVD Research" for non-DNA analysis, and for DNA analysis RES_DNA = "CVD Research" would be used?
Yes No

(This file ICTDER02 has been distributed to ARIC PIs, and contains the responses to consent updates related to stored sample use for research.)

8.a. Will the DNA data be used in this manuscript? Yes
 No

8.b. If yes, is the author aware that either DNA data distributed by the Coordinating Center must be used, or the file ICTDER02 must be used to exclude those with value RES_DNA = "No use/storage DNA"?
 Yes No

9. The lead author of this manuscript proposal has reviewed the list of existing ARIC Study manuscript proposals and has found no overlap between this proposal and previously approved manuscript proposals either published or still in active status. ARIC Investigators have access to the publications lists under the Study Members Area of the web site at: <http://www.csc.unc.edu/ARIC/search.php>

Yes No

