

ARIC Manuscript Proposal # 1759

PC Reviewed: 3/8/11
SC Reviewed: _____

Status: A
Status: _____

Priority: 2
Priority: _____

1.a. Full Title: Associations of traditional cardiovascular risk factors and high-sensitivity cardiac troponin T.

b. Abbreviated Title (Length 26 characters): Traditional risk factors and hs-cTnT.

2. Writing Group:

Jonathan Rubin, MD; Kunihiro Matsushita, MD, PhD; Christie M. Ballantyne, M.D.; Vijay Nambi, MD; Ron Hoogeveen, Ph.D.; A. Richey Sharrett MD, DrPH; Roger S. Blumenthal, MD; Josef Coresh, MD, PhD; Elizabeth Selvin, PhD, MPH; others welcome.

I, the first author, confirm that all the coauthors have given their approval for this manuscript proposal. JR **[please confirm with your initials electronically or in writing]**

First author: Jonathan Rubin, MD.
Address: Department of Epidemiology
Johns Hopkins Bloomberg School of Public Health
2024 E. Monument St., Room 1-500K
Baltimore, MD 21287
Phone: 410-612-9118 Fax: 410-955-0476
E-mail: jorubin@jhsph.edu

ARIC author to be contacted if there are questions about the manuscript and the first author does not respond or cannot be located (this must be an ARIC investigator).

Name: Elizabeth Selvin PhD, MPH
Address: Department of Epidemiology
Johns Hopkins Bloomberg School of Public Health
Welch Center for Prevention, Epidemiology, and Clinical Research
2024 E Monument Street, 2-600
Baltimore, MD 21287
Phone: 410-614-3752 Fax: 410-955-0476
E-mail: lselvin@jhsph.edu

3. Timeline: We aim to complete this manuscript within 6 months of approval.

4. **Rationale:**

Cardiac troponin-T (cTnT) measured in plasma allows for the noninvasive identification of persons at risk for a cardiovascular event ¹⁻⁵. A recently developed, high-sensitivity cardiac troponin-T (hs-cTnT) assay allows for the detection of hs-cTnT levels below clinical thresholds or detection levels of earlier cTnT assays. Using this novel pre-commercial assay, troponin levels far below the conventional limit of detection have been shown to predict cardiovascular events and death from any cause in persons with heart failure ⁶, stable coronary artery disease ⁷ and in the general population ^{8,9}. Unpublished data from ARIC (MS # 1563) suggests that hs-cTnT is also independently associated with heart failure and all-cause mortality.

Various mechanisms have been proposed to explain the release of troponin in persons with and without obstructive coronary heart disease including subendocardial ischemia leading to myocyte necrosis, oxidative stress and ischemia without necrosis ¹⁰. Other potential sources of elevated troponin include left ventricular strain ¹¹, apoptosis ¹², vigorous exercise ¹³ and cardiomyocyte turnover ¹⁴.

The determinants of elevated hs-cTnT levels below clinical thresholds in the general population are unknown. Measurement of hs-cTnT from stored samples from participants who attended the fourth ARIC visit provides a community-based sample in which to assess the relationship between traditional cardiovascular risk factors and hs-cTnT.

5. **Main Hypothesis/Study Questions:**

Aim: To characterize the cross-sectional associations between traditional cardiovascular risk factors and hs-cTnT

6. **Design and analysis (study design, inclusion/exclusion, outcome and other variables of interest with specific reference to the time of their collection, summary of data analysis, and any anticipated methodologic limitations or challenges if present).**

Study design: Cross-sectional analysis of participants who attended ARIC visit 4.

Exposure: Traditional cardiovascular risk factors (age, sex, diabetes status (undiagnosed, diagnosed, treated/untreated), race/center, smoking status, body mass index, waist-hip ratio, blood pressure, hypertensive medication use, education, income, triglycerides, total-, HDL- and LDL-cholesterol, metabolic syndrome, kidney function (estimated glomerular filtration rate (GFR) from serum creatinine) and Framingham Risk Score.

Outcome:
hs-cTnT modeled continuously and by categories (as described below in the Statistical Analysis section).

Inclusions

All black and white ARIC subjects with data on high-sensitivity troponin T measured from blood samples (plasma) obtained at visit 4 (n ≈ 11,500).

Exclusions:

Race other than black or white, missing hs-cTnT at visit 4, or missing covariates of interest.

Statistical Analysis

Continuous variables will be reported as means with standard deviations and categorical variables as proportions. We will model hs-cTnT as both a categorical and a continuous variable. For categorical analyses, Group 1 will include all participants with undetectable levels (~30%). The remaining (~70%) will be categorized into approximate thirds: hs-cTnT 0.003–0.005 µg/L (Group 2), 0.006–0.008 µg/L (Group 3), and higher levels further divided at approximately the 99th percentile value specified by the manufacturer (which corresponded to the ~90th percentile in the ARIC populations into Group 4 (0.009–0.013 µg/L) and Group 5 (≥0.014 µg/L). For continuous models, undetectable levels of hs-cTnT will be assigned a value of 0.0015 µg/L (i.e., half the lower limit of detection).

Statistical comparison of groups will be performed using one-way ANOVA for continuous variables or chi-square tests for categorical variables. Multivariable logistic regression models will be used to identify variables that are independently associated with elevated hs-cTnT. Analyses will be performed separately in persons with no previous clinical or silent CHD. We will also conduct analyses implementing linear splines and restricted cubic splines in multivariable models to characterize the shape of the association of traditional risk factors with hs-cTnT.

Limitations

Because neither echocardiography nor coronary angiographies were available from participants who attended ARIC visit 4, we cannot rule out elevated hs-cTnT due to underlying subclinical cardiovascular disease. Despite adjustment for known risk factors for cardiovascular disease, we will also not be able to rule out the possibility of residual confounding. The association with race cannot be easily separated from geography since the majority of blacks were recruited at the Jackson Field Center. Due to the cross-sectional nature of this investigation, the temporality of any observed associations cannot be established.

7.a. Will the data be used for non-CVD analysis in this manuscript?__ Yes X No

b. If Yes, is the author aware that the file ICTDER03 must be used to exclude persons with a value RES_OTH = “CVD Research” for non-DNA analysis, and for DNA analysis RES_DNA = “CVD Research” would be used?

 Yes No

(This file ICTDER03 has been distributed to ARIC PIs, and contains the responses to consent updates related to stored sample use for research.)

12. Manuscript preparation is expected to be completed in one to three years. If a manuscript is not submitted for ARIC review at the end of the 3-years from the date of the approval, the manuscript proposal will expire.

Reference

1. Melander O, Newton-Cheh C, Almgren P, Hedblad B, Berglund G, Engström G, Persson M, Smith JG, Magnusson M, Christensson A, Struck J, Morgenthaler NG, Bergmann A, Pencina MJ, Wang TJ. Novel and conventional biomarkers for prediction of incident cardiovascular events in the community. *JAMA*. 2009;302:49-57
2. Blankenberg S, Zeller T, Saarela O, Havulinna AS, Kee F, Tunstall-Pedoe H, Kuulasmaa K, Yarnell J, Schnabel RB, Wild PS, Munzel TF, Lackner KJ, Tiret L, Evans A, Salomaa V. Contribution of 30 biomarkers to 10-year cardiovascular risk estimation in 2 population cohorts: The monica, risk, genetics, archiving, and monograph (morgam) biomarker project. *Circulation*. 2010;121:2388-2397
3. Zethelius B, Berglund L, Sundström J, Ingelsson E, Basu S, Larsson A, Venge P, Arnlöv J. Use of multiple biomarkers to improve the prediction of death from cardiovascular causes. *N Engl J Med*. 2008;358:2107-2116
4. Lindahl B, Venge P, Wallentin L. Relation between troponin t and the risk of subsequent cardiac events in unstable coronary artery disease. The frisc study group. *Circulation*. 1996;93:1651-1657
5. Daniels LB, Laughlin GA, Clopton P, Maisel AS, Barrett-Connor E. Minimally elevated cardiac troponin t and elevated n-terminal pro-b-type natriuretic peptide predict mortality in older adults: Results from the rancho bernardo study. *J Am Coll Cardiol*. 2008;52:450-459
6. Latini R, Masson S, Anand IS, Missov E, Carlson M, Vago T, Angelici L, Barlera S, Parrinello G, Maggioni AP, Tognoni G, Cohn JN. Prognostic value of very low plasma concentrations of troponin t in patients with stable chronic heart failure. *Circulation*. 2007;116:1242-1249
7. Omland T, de Lemos JA, Sabatine MS, Christophi CA, Rice MM, Jablonski KA, Tjora S, Domanski MJ, Gersh BJ, Rouleau JL, Pfeffer MA, Braunwald E. A sensitive cardiac troponin t assay in stable coronary artery disease. *N Engl J Med*. 2009;361:2538-2547
8. deFilippi CR, de Lemos JA, Christenson RH, Gottdiener JS, Kop WJ, Zhan M, Seliger SL. Association of serial measures of cardiac troponin t using a sensitive assay with incident heart failure and cardiovascular mortality in older adults. *JAMA*. 2010;304:2494-2502
9. de Lemos JA, Drazner MH, Omland T, Ayers CR, Khera A, Rohatgi A, Hashim I, Berry JD, Das SR, Morrow DA, McGuire DK. Association of troponin t detected with a highly sensitive assay and cardiac structure and mortality risk in the general population. *JAMA*. 2010;304:2503-2512
10. Hickman PE, Potter JM, Aroney C, Koerbin G, Southcott E, Wu AH, Roberts MS. Cardiac troponin may be released by ischemia alone, without necrosis. *Clin Chim Acta*. 2010;411:318-323
11. Cheng W, Li B, Kajstura J, Li P, Wolin MS, Sonnenblick EH, Hintze TH, Olivetti G, Anversa P. Stretch-induced programmed myocyte cell death. *J Clin Invest*. 1995;96:2247-2259

12. Narula J, Haider N, Virmani R, DiSalvo TG, Kolodgie FD, Hajjar RJ, Schmidt U, Semigran MJ, Dec GW, Khaw BA. Apoptosis in myocytes in end-stage heart failure. *N Engl J Med*. 1996;335:1182-1189
13. Shave R, Baggish A, George K, Wood M, Scharhag J, Whyte G, Gaze D, Thompson PD. Exercise-induced cardiac troponin elevation: Evidence, mechanisms, and implications. *J Am Coll Cardiol*. 2010;56:169-176
14. Bergmann O, Bhardwaj RD, Bernard S, Zdunek S, Barnabe-Heider F, Walsh S, Zupicich J, Alkass K, Buchholz BA, Druid H, Jovinge S, Frisen J. Evidence for cardiomyocyte renewal in humans. *Science*. 2009;324:98-102