

ARIC MANUSCRIPT PROPOSAL FORM

Manuscript #621

1. **Full Title:** Does vital exhaustion increase the risk of stroke?

Abbreviated Title: (26 letters/space) Vital exhaustion and stroke

2. **Writing Group:**

Lead: Skai W. Schwartz, Ph.D.

Assistant Professor

Department of Epidemiology and Biostatistics

College of Public Health, MDC-56

13201 Bruce B. Downs Blvd

Tampa, FL 33612-3805

Phone: (813) 974-6679

Email: sschwartz@coml.med.usf.edu

Other authors: P. McGovern M.D., W. Rosamond Ph.D., E. Shahar M.D., L. E. Chambless Ph.D.

3. **Timeline:**

The analysis is contingent on there being at least 200 confirmed incident strokes or TIA's post visit 2. We understand that this is not expected until December, 1999. We hope the manuscript can be submitted very early in the next millennium.

4. **Rationale:**

The Maastricht questionnaire was developed to measure vital exhaustion, defined as "A state which is present when the individual not only complains of decreasing energy but also by feeling dejected or defeated.(1)" In addition to measuring feelings of helplessness, the questionnaire contains symptoms of insomnia (trouble falling asleep and waking up repeatedly during the night) and possible obstructive sleep apnea, a condition in which there is closure of the upper airway tract while the diaphragm continues its breathing effort. Apneic episodes lead to increased intracranial pressure and decreased cerebral perfusion pressure and can cause a marked reduction in cerebral blood flow in some patients(12-14). Exhaustion, obstructive apnea and subjectively reported sleep complaints such as "trouble falling asleep," have all been shown to be risk factors for coronary events (2-10, 15-21). It has been suggested that sleep complaints coupled with exhaustion represent an inability to cope with daily stressors(17) and recently daily stress has been correlated with myocardial ischemia(22). Although related symptoms such as excessive daytime sleepiness have been reported to precede stroke(11), vital exhaustion has not, to our knowledge, been studied in conjunction with stroke. However, McGovern et al. reported that in the ARIC cohort, while an age-adjusted measure of carotid wall thickness was related to excess fatigue, this relationship could be explained by antihypertensive medications, body mass index and conventional risk factors, particularly smoking (23). It is possible that excess fatigue, apnea and insomnia may affect stroke risk by a mechanism other than atherosclerosis.

5. **Main Issues/Hypotheses to be Addressed:**

(1) To determine the ability of the Maastricht questionnaire to predict stroke, after adjustment for conventional risk factors.

(2) To determine the items of the Maastricht questionnaire that are most predictive of stroke, in order to provide clues as to the mechanism.

Methods:

Vital exhaustion and stroke: A proportional hazards regression, using the date of onset of stroke symptoms will be the primary analysis used to determine if vital exhaustion increases the risk of stroke. Analyses will be done overall and by gender and by race.

Determining most predictive items: A logistic discriminant analysis will be used to rank items by predictive ability for stroke.

Power Considerations: If we assume a population of 12,000 individuals with 230 incident strokes, then if 20% of the subjects are considered exhausted, there is about 95% power to detect a risk ratio of 1.75 or higher. The table below varies the prevalence of VE and the risk ratio detected.

Table: Statistical Power Estimates for the association of vital exhaustion and incident stroke, assuming an average sample size of 12,000 persons at risk and 230 incident strokes.

<u>Prevalence of</u> <u>exhaustion</u>	<u>Risk ratio</u>	<u>Power</u>
10% (e.g.: upper decile of questionnaire)	1.5	57%
	1.75	84%
20% (e.g.: upper quintile)	1.5	74%
	1.75	95%
25% (e.g.: upper quartile)	1.5	80%
	1.75	97%

6. Data:

ARIC Data for Visit 2 and later will be required.

Visit 2 data required:

Visit 2 date
 Individual Items of the Maastricht questionnaire and
 Total score on the Maastricht questionnaire
 Previous/current Stroke indicator
 Previous/current TIA indicator
 Study center
 Gender
 Race
 Age
 Education in years
 Body mass index (kg/m²)
 Smoking category: Current, Former or Never smoker
 Diabetes mellitus indicator
 Systolic blood pressure
 Total Cholesterol
 HDL Cholesterol
 Triglycerides
 Fibrinogen

All Visits post Visit 2

Stroke indicator

TIA indicator
Stroke onset date
TIA onset date
1st stroke versus not

Also characteristics of the strokes or TIAs for a sentence in the test
1st stroke versus not
Ischemic or other
Right of Left side

References

1. Appels A, Hoppener P, Mulder P: A Questionnaire to Assess Premonitory Symptoms of Myocardial Infarction. *International Journal of Cardiology* 1987; 17; 15-24
2. Alonzo A, Simon A, Feinleib M: Prodromata of Myocardial Infarction and Sudden Death. 1975 *Circulation*, 52, 1056-1062
3. Feinleib M, Simon AB, Gullum RF, Margolis JR: Prodromal Symptoms and Signs of Sudden Death. Supplement III to *Circulation*, Vols 51 and 52, December 1975 III-155-159
4. Greene SW, Goldstein S, Moss AJ: Psychosocial Aspects of Sudden Death. *Arch Intern Med* 1972; 129: 725-731
5. Kuller L: Prodromata of Sudden Death and Myocardial Infarction. *Adv Cardiol* 25: 61-72; 1978
6. Kaye J, Kaye K, Madow L: Sleep Patterns in Patients with Cancer and Patients with Cardiac Disease. *Journal of Psychology* 1983; 114, 107-113
7. Romo: Factors Related to Sudden Death in Acute Ischemic Heart Disease. *Acta Med Scand*. 1973; 547: 5-9
8. Master AM, Dack S, Jaffe HL: Premonitory Symptoms of Acute Coronary Occlusion: A Study of 260 Cases. *Ann Intern Med* 14: 1155, 1941
9. Stowers M, Short D: Warning Symptoms Before Major Myocardial Infarction. *Br Heart J*, 32: 833, 1970
10. Nixon PGF, Bethel HJN: Preinfarction Ill Health. *Am J Cardiol* 33: 446, 1974
11. Van Diest R: Subjective Sleep Characteristics as Coronary Risk Factors, Their Association with Type A Behavior and Vital Exhaustion. *J of Psychosomatic Research* 34(4):415-426, 1990
12. Palomaki H, Partinen M, Ertinijuntti T, Kaste M: Snoring, sleep apnea syndrome and stroke. *Neurology* 1992; 42(suppl 6): 75-82
13. Bassetti C, Aldrich MS, Chervin RD, Quint D: Sleep apnea in patients with ischemic attack and stroke; *Neurology* 47(5):1167-73 1996 Nov.
14. Pressman MR, Schetman WR, Figueroa WG: Transient ischemic attack and minor stroke during sleep: relationship to obstructive sleep apnea syndrome. *Stroke* 26(12):2361-5, 1995 Dec.
15. Friedman G, Ury H, Klatsky A, Siegelbaum A: A Psychological Questionnaire Predictive of Myocardial Infarction: Results From the Kaiser Permanente Epidemiologic Study of Myocardial Infarction.

16. Eaker ED, Pinsky J, Castelli WP: Myocardial Infarction and Coronary Death Among Women: Psychosocial Predictors From a 20 Year Followup of Women in the Framingham Study. *Am J Epidemiol* 1992;135:854-64
17. Siegrist J, Peter R, Motz W, Strauer BE: The Role of Hypertension, Left Ventricular Hypertrophy and Psychosocial Risks in Cardiovascular Disease: Prospective Evidence from Blue Collar Men. *European Heart J* 1992; 13(suppl D), 89-95
18. Koskenvuo M, Kaprio J, Partinen M, Langinvainio H, Sarna S, Rita H, Heikkila K: Poor Sleep Quality, emotional Stress and Morbidity: A six Year Followup of 10778 persons aged 35-59 years. In *Stress and Psychosomatics*(edited by Achte K, Pakaslahti A) pp 115-20 Helsinki Psychiat Fennica (Suppl) 1986
19. Appels A, De Vos Y, Van Diest R, Hoppener P, Mulder P, De Groen J: Are Sleep Complaints Predictive of Future Myocardial Infarction? *Activitas Nervosa Super (Praha)* 1987; 29: 147-151
20. Vanhalla, K. Sepelvaltimotautien psyykkisistä riskitekijöistä. Prospektiivinen Tutkimus helsinkiläisten poliisimiesten sepelvaltimotautiin sairautuvuutta ennustavista psyykkisistä tekijöistä. [Psychological Risk Factors Related to Coronary Heart Disease Morbidity of Helsinki Policemen]. Monographs of Psychiatria Fennica. Editor: K.A. Achte, Chief and Professor of Psychiatry, Psychiatric Clinic, Helsinki University of Central Hospital, Lapinlahdentie, 00180 Helsinki 18, Finland. Printed in Kyriiri, Finland, 1979, ISBN 951-709-041-2 ISSN 0355-7707
21. Schwartz SW. Sleep Complaints As A Risk Factor For Coronary Heart Disease Events in Older Adults: Evidence from the Piedmont Study. Doctoral Dissertation: University of North Carolina at Chapel Hill, 1995.
22. Guille EC, Blumenthal JA, Babyak M, Jiang W, Waugh RA, Frid DJ, O'Connor CM, Morris JJ and Kranz DS: Effects of Mental stress on Myocardial ischemia during daily life. *JAMA*, 1997;277(20):1521-1526
23. McGovern PG, Shahar E, Folsom AR, Rosamond WD: No relation between excess fatigue and asymptomatic carotid sclerosis. *Epidemiology* 1996;7:638-640.